

MESSENGER-INQUIRER



October 21, 2023

Cleaning Up the Garden Reduces Insect Pests and Plant Diseases for the Next Season

A way to manage and reduce insect pests in the garden for next year is to remove weeds and diseased plant debris this fall. By removing finished vegetable plants and dried fruit from trees, the amount of insect pests and disease organisms are reduced at the beginning of the next growing season in the garden.

Insect pests may be found in dead leaves in and around the garden. Pests such as squash vine borer, Mexican bean beetle, squash bug, diamondback moth, tomato hornworm, cabbage looper, and imported cabbageworm are a few examples of insects able to overwinter in the garden. Getting rid of the dead tomato vines, bean bushes, squash plants, cabbage plants, and other plant material lessens the chance that these pests will overwinter in the garden. These insects often make their home in dead leaves and plant material, which offers them some protection from the elements. Removing the plants from the area of the vegetable garden or tilling them into the soil destroys their hiding places.

MESSENGER-INQUIRER

Many diseases overwinter on the same plants they infected during the summer. For example, apple scab, a common fungal disease that infects apples and crab apples, spends the winter in dead infected leaves on the ground. The warm, moist weather of spring triggers the fungus to grow and release spores that infect new growth on the tree. Raking up dead leaves from under these trees in the fall is one way to reduce the severity of disease next spring. The disease may not be completely controlled by this method because of the weather, but good sanitation practices are an important part of any disease control strategy.

Get rid of rotting fruit from trees, especially fruit that is hanging on the tree. Even though this fruit seems dry and lifeless, it is likely to harbor many diseases. This is also true for grapes. During the pruning process in late winter, remove fruit that is still on the vine, which may serve as a source of disease.

Practicing good sanitation in the vegetable garden is important too. Vegetable plant tissues can harbor diseases over the winter. Remove dead tomato plants, cucumber vines, squash plants, and other plant material. If the material is destroyed, then it will not be an immediate source of disease for the next year and slows down the development of common plant diseases.

Excessive weed growth during late fall and winter also contributes to pest problems. Weeds provide food and shelter for many overwintering insects and serve as sites for egg laying in the fall. Flea beetles, which are pests on young vegetables, especially sweet corn and eggplant, find food and shelter in crop residues and weeds throughout the winter.

MESSENGER-INQUIRER

The two-spotted spider mites continue to feed on weeds after the vegetable crops have withered. If the weeds are not removed or mowed down, the spider mites will be ready to infest your vegetables next spring.

In the landscape, twig girdlers have been active in hickory, pecan, and persimmons. The female beetle selects a twig about the diameter of a fat pencil and chews deep, narrow grooves that leave about a two-foot-long section attached by only a slender piece of heartwood. The brown beetles crawl along the terminal portion and make small notches in which 5 to 20 eggs are placed. The girdled twigs, which contain the eggs and white legless larvae, break and fall to the ground. The twig pruning produces growth deformities that affect the shape and appearance of small trees.

To manage the twig girdlers, collect and destroy the fallen twigs. This is the most effective means for reducing the potential infestation for next year. Application of insecticides to control these insects has not been very satisfactory.

Another common landscape pest is the bagworm. The small, 1.5- to 2-inch long, brown bags hanging in evergreen trees and shrubs are full of several hundred eggs. The eggs will hatch in May. By removing the bags now, their population will be reduced. The easiest way to remove the bag is with hand pruners or heavy scissors, otherwise, the needles or scales of the evergreen may be stripped off the branch as the bag is pulled. If there are too many to remove by hand, watch for the eggs to hatch in mid-May to June. The small larvae are easier to manage with an insecticide at that time.

MESSENGER-INQUIRER

Fall and winter are also excellent times to inspect plants for scale insects. These insects usually have a hard outer covering similar to a shell and are found where branches join the trunk or near buds. Scale insects may also be seen this time of year on the leaves of broadleaf evergreens such as holly or euonymus. Treatments with dormant or horticultural oils are usually effective against scale. Follow label directions when applying insecticides.

For more information about cleaning up the garden and landscape, contact the Daviess County Cooperative Extension Service at 270-685-8480 or annette.heisdorffer@uky.edu.

Annette's Tip:

Leave hummingbird feeders out because the stragglers migrating south need food along the way. Leaving the feeder out does not keep them from migrating. Hopefully, most have moved on with the cooler weather arriving.

Ongoing Event:

The Owensboro Regional Farmers Market is still open on Saturdays from 8:00 a.m. to noon through November 4. A special holiday market will be held on November 4 with extra vendors including crafts for the holidays.

Good News and Congratulations:

Congratulations to Dr. Cathy Schmitt for being named Outstanding Master Gardener Individual 2023 at the Extension Master Gardener State Conference recently held in Frankfort, KY.

Educational programs of Kentucky Cooperative Extension serve all people regardless of economic or social status and will not discriminate on the basis of race, color, ethnic origin, national origin, creed, religion, political belief, sex, sexual orientation, gender identity, gender expression, pregnancy, marital status, genetic information, age, veteran status, or physical or

MESSENGER-INQUIRER

mental disability. UNIVERSITY OF KENTUCKY, KENTUCKY STATE UNIVERSITY, U.S.
DEPARTMENT OF AGRICULTURE, AND KENTUCKY COUNTIES, COOPERATING