

July 12, 2025

Chiggers Spoil Summertime Fun By Annette Meyer Heisdorffer, PhD Daviess County Extension Agent for Horticulture Education

Enjoying the outdoors and ending up with red spots on the ankles and in thin skin areas on the body is frustrating. Chiggers may be the culprit to blame. Information about their preferred habits, life cycle, and some management tips are provided in this article.

Chiggers are actually immature mites. They cause an aggravating rash and intense, itchy feeling when they feed on the skin of animals and humans, according to Dr. Jonathan Larson, University of Kentucky Extension Entomologist. Chiggers are not seen in the landscape or on your body because they are so tiny. Under the microscope, they can be seen with six legs and reddish color.

Chiggers that attack people are the immature form of a predatory mite as an adult. During the immature larval stage, they digest skin cells and use them for food. They pierce the skin with their tube-like mouthpart, inject "saliva" which destroys the skin cells, and consume the digested material. They can continue this process over 2 to 4 days. A misconception is that the chiggers burrow into the skin, but this is not correct. They stay as an external parasite on top of the skin.

Chiggers usually feed in areas where the skin is thinner or where the area is constricted by clothing. This is why common feeding sites are the ankles, behind the knees, the groin area, the waistband area, in armpits, and around bra lines, according to Dr. Larson.

It may be 12 to 24 hours after the chigger feeds before noticing the bitten area. This inflamed area becomes reddish in color. At the center of this area, it may be sunken with a red dot or red and slightly raised, and possibly a pustule forms, acting like a blister when popped. Everyone reacts differently. Chigger bites can be treated with over-the-counter medications such as hydrocortisone or calamine lotion. Consult a medical professional for more help.

What is the life cycle of chiggers? The adult stage spends the winter in the soil. When the soil temperature reaches around 60 degrees F, the adults emerge and mate. The female lays her eggs in the soil in areas that are overgrown. The eggs hatch and the chiggers, crawl up the grass or on other plants and wait for a host such as a human. When finished feeding, the chigger drops off and molts. There are multiple generations of chiggers with populations peaking in the spring and late summer.

How can we protect ourselves from chiggers? It is best to avoid unmown areas or areas that are overgrown. Try to stick to trails that have been managed. When needing to work in overgrown areas, wear long pants and tuck them into socks or shoes. Also, wear long sleeves. When finished and coming inside from working outside, wipe down your legs with a paper towel or take a shower, and especially scrub your legs and ankles.

If you want to use an insect repellent on your skin, use those that include DEET, picaridin, IR-3535, or oil of eucalyptus. These would also be used to repel ticks and mosquitoes.

Multiple applications may be needed depending on the product. Consult the label for more information on how the product can be used.

Another option is to treat your clothes with permethrin. Use this product according to the label. It is only to treat clothing before the clothing is worn because it is a true insecticide and should not be placed on the skin. Always read and follow label directions for the safe use of pesticide. Chiggers, ticks, and other pests will die when they climb onto the permethrin treated clothes.

To discourage habitats for chiggers, remove brush, mow, and manage the landscape. An increase in light and a reduction in humidity decreases the attraction of the area to the chiggers.

If there is a hot spot of chiggers, this area could be treated with permethrin, bifenthrin, or carbaryl for about 1 to 2 weeks of control depending on weather conditions and the product. Follow the label directions. This is certainly not a long-term solution. Treatment needs to start in April or May. A serious infestation may need another application in June.

For more information about chiggers and how to manage them, contact the Daviess County Cooperative Extension Service Office at 270-685-8480 or annette.heisdorffer@uky.edu. Annette's Tip:

Slime molds may appear in 4- to 6-inch patches of white, cream, gray, or purple with a crusty surface, looking like dog vomit, in the landscape. They do not hurt plants but use them as support during their reproductive state.

Upcoming Event:

The 2025 Extension Master Gardener program will be offered this fall. An application and interview are required. This 14-week program will be held on Thursdays, from 9:00 a.m. to

noon, starting September 4. This is a volunteer program with a requirement of 40 hours given back through approved Cooperative Extension Service programs. The training includes presentations, quizzes, and an open-book test. Trained Extension Master Gardeners help the University of Kentucky Martin-Gatton College of Agriculture, Food and Environment Cooperative Extension Service by educating the public with science based horticultural information to protect the environment as well as to use proper techniques to manage plants for the landscape and food production. The cost is \$100. Applications are available and must be returned to the office by July 25. Contact the Daviess County Extension Office if interested in completing this volunteer program.

The Martin-Gatton College of Agriculture, Food and Environment is an Equal Opportunity Organization with respect to education and employment and authorization to provide research, education information and other services only to individuals and institutions that function without regard to economic or social status and will not discriminate on the basis of race, color, ethnic origin, national origin, creed, religion, political belief, sex, sexual orientation, gender identity, gender expression, pregnancy, marital status, genetic information, age, veteran status, physical or mental disability or reprisal or retaliation for prior civil rights activity.